## TUBE FEEDING THE HYPOACTIVE PATIENT

- 1. Explain the procedure to the patient stressing the need to maintain his nutritional status
- 2. Provide privacy and send prepared feeding tray to patient's bedside
- 3. Put patient in the most comfortable position, together with the assistant
- 4. Assign the assistant to study the patient's head
- 5. Wash, dry hands and check if the NG tube is in stomach
- 6. Remove spigot and fix the barrel of 20cc syringe to the NG tube
- 7. Run about 10-15 mils of normal saline or water through the tube
- 8. Check temperature of feed and pour it into syringe and allow to run by gravity
- 9. Pour the feed gradually until the required amount is administered
- 10. Observe the patient throughout the procedure
- 11. Run water(10-15mils) through the NG tube after feeding
- 12. Remove the syringe, clean the tip of the tube and fix the spigot and secure the NG on the patient's neck
- 13. Make patient comfortable and thank him
- 14. Discard tray and wash hands

Name of staff(MO/NO/CO/COI/MIDWIFE/ETC)

15. Document procedure on the intake and output chart and nurse's note indicating patient's reaction and any other observations

## **SIGNATORIES**

Name: Department in charge	sign	Date:
Name: Nursing Officer in charge	sign	Date:
Name:	sign	Date: